Description of a new Species of Pseudacræa in the Rev. Mr. CORNTHWAITE'S *Collection:* by A. G. BUTLER, F.L.S., F.Z.S., &c.

GENUS PSEUDACRÆA. *Westwood.*

Pseudacræa simulator, n. sp.

♀. *Wings above* olive-brown with the nervures and intermediate folds black; frontwings with basal area blackish; five black spots at base, three within cell and two below it, two of them touching base, that within cell having a white pupil, all of them with a griseous iris; a broad white band nearly equal in width throughout (interrupted by the first and second subcostal branches, the radials, upper disco-cellular, and second and third median branches), crosses the centre of the wing obliquely from costal nervure to below centre of first median interspace; three white rays on inner margin: *hindwings* with base tawny; eleven black spots, the two touching the base being largest and having conspicuous white pupils; a broad externally diffused white band crosses the wing from inner margin to costal nervure, and is about twice as broad at its lower as at its upper extremity: body moderately robust, thorax black, white-spotted, abdomen bright ochre-yellow, with diffused dorsal black stripe; antennæ black, palpi wanting, but apparently bright ochre-yellow.*

Wings below paler than above; an additional black spot towards end of frontwing cell; otherwise as above: expanse of wings, 3 inches, 4 lines.

West Africa. Coll. Rev. T. Cornthwaite.

This species is a remarkable mimic of *Acræa (Planema) Euritea* ♀; it is allied to *P. Imitator* of Trimen.

---

* A remnant of one of them remains sticking to the proboscis.